

THE ISOPERIMETRIC INEQUALITY VIA APPROXIMATION THEORY AND FREE BOUNDARY PROBLEMS

CATHERINE BÉNÉTEAU AND DMITRY KHAVINSON

Abstract. In this survey paper, we examine the isoperimetric inequality

One of the early analytic proofs of (1.1) was given by A. Hurwitz [26] in 1901. Let us sketch his argument here.

Sketch of proof. Suppose the region Ω is bounded by the simple closed smooth curve γ ; parametrized with respect to the arc-length parameter s and with length 2ℓ : (So, the isoperimetric inequality would state that $A \leq \ell^2$).

problems. This approach reveals a close tie to hydrodynamics and, in particular, to problems concerning shapes of electrified droplets of perfectly conducting fluid. We use as a point of departure the paper [32], in which the author discusses the concept of analytic content, and the related survey paper [18]. In Section 3, we discuss the connection with overdetermined boundary value problems and Serrin's theorem. In Section 4, we describe a more general problem and its application to determining the shape of a droplet of conducting fluid in the presence

This theorem is discussed in detail in [18]. Let us outline the argument here. H. Alexander in 1973 ([3]) proved the upper estimate by noticing the connection with the Ahlfors-Beurling estimate from 1950 ([1]).

More specifically, suppose D is a bounded domain (with smooth boundary ∂D) containing z_1 : By the Cauchy-Green formula,

$$\mathfrak{J} = \frac{1}{2\pi i} \int_{\partial D} \frac{z}{z_1 - z} dz + \frac{1}{\pi} \int_D \frac{1}{z_1 - z} dA(z);$$

where dA is area measure. Define

$$G(z) = \frac{1}{\pi} \int_D \frac{1}{z_1 - z} dA(z);$$

Then

$$\mathfrak{J} + G(z) = \frac{1}{2\pi i} \int_{\partial D} \frac{z}{z_1 - z} dz + 1$$

Theorem 2.2. ([32]) *Let Ω and γ be as above. The following are equivalent:*

(i) $\chi_\Omega = \frac{2A}{P}$;

(ii) *There is $\gamma \in A_\Omega$ such that $\dot{z}(s) \perp \dot{\gamma}(s) = \dot{z}(s)$ on γ ; where s is the arc-length parameter;*

(iii) $\frac{1}{A} \int_\Omega f dA = \frac{1}{P} \int_\gamma f ds$ for all $f \in A_\Omega$;

Remark. Note that (iii) holds for annuli $\Omega = \{r < |z| < R\}$. Simply take the Laurent series decomposition of $f = f_1 + f_2$ in the annulus, where f_1 is analytic inside Ω : $|z| < R$ and $f_2(1) = 0$; and notice that both sides of the equality in (iii)

The left hand side of equation (2.3) is real and therefore has argument increment 0; while the right hand side has an argument increment of at least 4π as we travel along γ ; unless γ is a closed curve.

Indeed, take any harmonic test function u in Ω - that is smooth up to the boundary. By Green's formula,

$$\int_{\partial\Omega} uv_n ds = \int_{\Omega} \Delta u dx = \frac{A}{P} \int_{\partial\Omega} u ds:$$

Since u is arbitrary, $v_n = A/P$ on $\partial\Omega$: In this context, the shape of Ω was already known. We state the following result due to Serrin in two dimensions, although the theorem is more general and holds in all dimensions.

Theorem 3.2. ([56]) *If the overdetermined boundary value problem*

$$\Delta v = 1 \text{ in } \Omega;$$

$$v = 0 \text{ on } \partial\Omega;$$

$$v_n = \text{const on } \partial\Omega;$$

has a smooth solution in Ω ; then Ω is a disk.

This leads to an equivalent form of Conjecture 2.1 "à la Serrin" ([35]):

Conjecture 3.1. *Let Ω be a multiply connected domain. If the overdetermined boundary value problem ($n \geq 2$)*

$$\Delta v = 1 \text{ in } \Omega$$

$$\frac{\partial v}{\partial n} = \frac{A}{P}$$

where p is the pressure at each point $(x; y; z)$ and depends only on z ; since the flow is laminary. Since $\frac{d^2 p}{dz^2} = 0$; the right hand side is actually a constant A ; giving us (modulo a constant multiple) the first equation in Serrin's theorem.

On the other hand, the force exerted by the water on the walls of the pipe is given by

$$F = (p_j \frac{4}{3} \tau \circ \psi)$$

We now turn to a discussion of condition (ii) in Theorem 2.2 and a related application to determining the shape of droplets of conducting fluid in the presence of an electric field.

4. Droplets

Recall one of the equivalent conditions for $\mathcal{J}_\lambda(-) = 2A=P$:

$$\dot{z}(s) \perp i_s \dot{z}(s) = \lambda(z(s))$$

for some $\lambda \in \mathbb{R}$: We would like to consider a more general problem, in which the function λ may not be continuous in the closure of Ω and may possibly have

exists by Koebe's theorem (see [20, p. 237-238]). Since Ω has a rectifiable boundary $\partial\Omega$, f'' can be shown to be in $E^1(\Omega)$: We say that Ω is a *Smirnov domain* if, for each $z \in \Omega$,

$$\log |f''(z)| = \frac{1}{2\pi} \int_{\partial\Omega} \log |f''(\zeta)| g_{\Omega}(z, \zeta) d\mu(\zeta)$$

Theorem 4.2. ([36, p. 24-26]) *There exists a one parameter family of unbounded domains Ω_t ; each with rectifiable boundary Γ_t ; and a corresponding family of functions F_t analytic in Ω_t except for a simple pole with residue 1 at $z=1$; such that*

$$F_t(z) = p_t z + i \zeta_t \frac{dz}{ds} \text{ on } \Gamma_t;$$

for some real constants p_t and ζ_t ; with $p_t \neq 0$; $\zeta_t \neq 0$:

Each of these domains Ω_t is thus an example of a solution to Problem 4.1. Their boundaries Γ_t are images of the unit circle under a rational mapping of degree 3 on which (4.1) holds. None of these curves however is a physical droplet. To our knowledge, no other examples of such domains are known. In particular, we do not know of any examples of transcendental curves satisfying (4.1), although, most likely, there are plenty of them!

Applying electrical forces to droplets of conducting fluid has led to some very concrete applications: the process of "electrowetting", for example, in which an electric force is applied at the interface of a droplet of conducting fluid and a solid, has applications to digital cameras, camera phones, and home security systems. In 2003, scientists from Philips Research created a fluid lens that operates on the basis of the process of electrowetting: two non-mixing fluids, one conducting and one not, are placed inside a tube. The layer between the liquids (the *meniscus*) acts as a lens. An electric field is applied to the tube, which causes the conducting fluid to change its shape, thus resulting in a change of the focal length of the lens. See [49] for more details. For further references on electrowetting and its applications, see [5, 24]. A slightly different type of application can be found in [11]: there, the authors use Schwarz functions to model the changing shape of a void created and traveling inside a thin metal conductor subjected to an intense electric field. This model is similar in some ways to the one used for Hele-Shaw flows (see [10, 23, 53]).

5. Some special cases

Let us now examine three distinguished cases of Problem 4.1, in which the boundary condition on $\Gamma = \bigcup_{j=1}^n \Gamma_j$ simplifies to one of the following:

$$(5.1) \quad F(z) = p_j z \quad z \in \Gamma_j; p_j \in \mathbb{R}; p_j \neq 0;$$

$$(5.2) \quad F(z) = i \zeta_j \frac{dz}{ds} \quad z \in \Gamma_j; \zeta_j \in \mathbb{R}; \zeta_j \neq 0;$$

$$(5.3) \quad F(z) = p_j z + c_j \quad z \in \Gamma_j; p_j \in \mathbb{R}; p_j \neq 0; c_j \in \mathbb{C};$$

Note that the existence of a function F satisfying (5.2) implies the existence of a function g satisfying (5.3): simply define

$$g(z) = \int (F(z))^2 dz.$$

Then, by (5.2), for $z \in \gamma_j$; we have

$$\int (F(z))^2 dz = i \int_j^2 \left(\frac{dz}{ds} \right)^2 dz = i \int_j^2 (z + c_j);$$

for some constant c_j . Therefore g is well-defined as a single valued analytic function, and (5.3) holds. From now on, we shall always assume additional regularity for Ω , i.e., that Ω is a Jordan Smirnov domain.

5.1. The Schwarz function.

generality $p_j = 1$), then Ω is a so-called *quadrature domain*; namely, if f is any function analytic in Ω ; then by the complex form of Green's theorem,

$$\int_{\Omega} f dA = \frac{1}{2i} \int_{\partial\Omega} f \bar{z} dz = \frac{1}{2i} \int_{\partial\Omega} f F dz = \sum_{j=1}^n f(a_j) \text{Res}_{a_j} F.$$

These domains have been intensely studied in the 1980s by D. Aharonov, B. Gustafsson, H. S. Shapiro, K. Ullemar, Y. Avsi (see [57] and references therein). Also, see [23] for an account of many recent developments.

Even when we do not require the coefficients p_j to be equal, a similar argument as in the proof of Theorem 5.1 shows that if Ω is assumed to have a simple pole at the origin and Ω is bounded, then Ω must be a disk. (This is well-known in the context of Schwarz functions: the Schwarz function of a domain has one pole if and only if the domain is a disk.) If the function F has two different poles (and if the coefficients p_j are different), then the problem is already more difficult.

5.2. **Vekua's Problem.** The second special case (5.2)

$$F(z) = \sum_{j=1}^n \frac{c_j}{z - a_j} \quad z \in \Omega; \quad \sum_{j=1}^n c_j = 0$$

is a particular example of an overdetermined boundary value problem made enormously popular by works of I. N. Vekua in the 1950s. It is not difficult to

holds in all dimensions provided that γ is a C^2 surface (see [52]). Equivalently, if the overdetermined boundary value problem

$$\begin{aligned} \Delta u &= 0 \text{ in } \Omega; \\ u &= \text{const} \neq 0 \text{ on } \gamma; \\ \frac{\partial u}{\partial n} &= \text{const on } \gamma \end{aligned}$$

has a solution in an (unbounded!) domain Ω , then γ is a circle.

Remark. In [14], the authors notice that it is possible to drop the assumption that the domain is Smirnov, but then instead one must assume that the function F is in E^2 ; since the proof uses the fact that the function $z^2(F(\phi(z)))^2 \phi'(z)$ is in $H^1(D)$ (where ϕ is the Riemann mapping from the disk to Ω), and therefore cannot coincide with the conjugate of an H^1 function on the circle. It is not clear whether the theorem itself fails if one drops the assumption that Ω is Smirnov and considers only $F \in E^1$: In this context, one must cautiously observe that in non-Smirnov domains, there exist functions with positive and bounded boundary values which belong to any E^p class, $p < 1$ (see [30]).

We may also consider the case where F has a simple pole at infinity. Recall that this context has a physical interpretation, discussed in Section 4, as a droplet of conducting fluid in which the surface tension is much larger than the pressure inside the droplet (which is then considered negligible). In this case, the following theorem gives an example of a family of mathematical droplets.

Theorem 5.4. ([36, Thm 6.2]) *Let γ be a Jordan curve, with (logarithmic) capacity 1; whose exterior Ω is a Smirnov domain. If $\zeta \geq \frac{3+2\sqrt{3}}{3}$ and there exists $F \in E^1$ near the boundary of Ω and with a simple pole at ∞ ; that is, $F = z + O(\frac{1}{z})$; and*

$$(5.5) \quad F = i\zeta \frac{dz}{ds} \text{ on } \gamma;$$

then γ is included into one parameter family f_t ; $t = 1/\zeta$, where f_t is the image of the unit circle under the conformal mapping

$$f_t(w) = \frac{1}{w} + 2tw + \frac{t^2}{3}w^3.$$

For $\zeta \geq \frac{3+2\sqrt{3}}{3}$; (5.5) has no solution among mathematical droplets with Jordan boundaries. The droplets are convex for $\zeta \geq 3$ and the family contains only one physical droplet corresponding to the value $\zeta = 3$.

6. Extensions to higher dimensions

Finally, let us discuss what is known in higher dimensions. Suppose Ω is a bounded domain in R^n ; $n \geq 3$; γ is the boundary of Ω ; V is the volume of Ω ; and P is the (surface) area of γ : Let $H(\Omega)$ be the closure in the uniform norm on $\bar{\Omega}$ of the space of functions harmonic in a neighborhood of Ω : More generally, if K is a compact subset of R^n ; and $C(K)$ is the space of continuous functions on

K ; we will write $H(K)$ for the uniform closure in $C(K)$ of the space of functions harmonic in a neighborhood of K :

Let $\mathbf{x} = (x_1, \dots, x_n)$ be a vector in \mathbb{R}^n ; and $|\mathbf{x}|^2 = \left(\sum_{j=1}^n x_j^2\right)^{\frac{1}{2}}$: If one thinks of $H(K)$ as the uniform closure of the kernel of the Laplace operator Δ and $R(K)$ as the uniform closure of the kernel of the operator $\Delta - \Delta_{\mathbf{x}}$; then the analogy of the anti-analytic function \bar{z} is the function $|\mathbf{x}|^2$; since $(\Delta - \Delta_{\mathbf{x}})(|\mathbf{x}|^2) = 1$ and $\Delta(|\mathbf{x}|^2) = 2n = \text{const} \neq 0$: With this in mind, we define the concept of *harmonic content* as follows.

Definition. *The harmonic content of K is defined to be*

$$\alpha(K) := \text{dist}_{C(K)}(|\mathbf{x}|^2; H(K)):$$

For a bounded domain Ω ; we will write $\alpha(\Omega) := \alpha(\bar{\Omega})$: We then have the following result.

Theorem 6.1. ([33])

$$\alpha(K) = 0, \quad H(K) = C(K):$$

Note that in the case of analytic content in \mathbb{C} ; the equivalence of the statements $\alpha(K) = 0$ and $R(K) = C(K)$ follows at once from the Stone-Weierstrass theorem, since $R(K)$ is an algebra. However, Theorem 6.1 is non-trivial, since $H(K)$ is not an algebra. Different proofs were given by Poletsky ([50]) and Bliedtner (see [7] and references therein, in particular to the works of W. Hansen).

Harmonic content can be estimated in terms of geometric quantities. If R_{harm} is the radius of the ball with the same capacity as Ω ; and R_{vol} is the radius of the ball with the same volume as Ω , then the following theorem gives upper and lower bounds for the harmonic content of a domain Ω :

Theorem 6.2. ([33, 34])

$$\frac{1}{2}R_{\text{harm}}^2 \cdot \alpha(\Omega) \leq \frac{1}{2}R_{\text{vol}}^2$$

and equality on either side occurs only for balls.

The upper estimate was proved in [33], and the lower estimate as well as extensions of both inequalities to general elliptic operators were obtained in [34]. An interesting extension of this result to approximation in C^1 -norm by harmonic

Recall that analytic content for a domain Ω in \mathbb{C} is defined as

$$A(\Omega) := \inf_{f \in \mathcal{A}(\Omega)} \int_{\Omega} |f_z|^2 dx dy$$

Note that this is also equal to

$$A(\Omega) := \inf_{f \in \mathcal{A}(\Omega)} \int_{\Omega} |f_{\bar{z}}|^2 dx dy$$

An anti-analytic function $f = f_1 + if_2$ can be identified with the harmonic vector field $\vec{f} = (f_1; f_2) = \nabla u$; u a harmonic real-valued function, where

$$\text{Div} \vec{f} = \text{Curl} \vec{f} = 0:$$

the extremal solids are *not* balls in all dimensions $n \geq 3$; although they are very symmetric algebraic surfaces that are getting more and more tightly sealed to the tangent plane at the maximum point (see [22, p. 82]). The following conjecture proposed in [22] remains open.

Conjecture 6.1. $\mu_1(\Omega) = R_{vol}$.

The following theorem is the analogue of Theorem 2.2 and gives conditions equivalent to the attainment of the lower bound in Theorem 6.3.

Theorem 6.4. ([22]) *TFAE*:

(i) $\mu_1(\Omega) = \frac{nV}{P}$:

(ii) There exists $f \in C^1(\bar{\Omega})$: $f|_{\partial\Omega} = 1$; $\mu_1(\Omega) = \int_{\Omega} |\nabla f|^2 dx$ on Ω ; where ν is the outward unit normal to $\partial\Omega$.

(iii) $\int_{\Omega} \frac{1}{V} u dV = \frac{1}{P} \int_{\partial\Omega} u d\mathcal{H}^n$ for all u harmonic in Ω such that $\int_S \frac{\partial u}{\partial n} d\mathcal{H}^n = 0$ for all closed surfaces S in Ω .

(iv) There exists u in Ω satisfying

$$\Delta u = 1;$$

$$\frac{\partial u}{\partial n} \Big|_{\partial\Omega} = \text{const};$$

$$u|_{\partial\Omega} = \text{local constant};$$

The following conjecture thus follows naturally:

Conjecture 6.2. $\mu_1(\Omega) = \frac{nV}{P}$, Ω is either a ball or a spherical shell.

Serrin's theorem in higher dimensions implies that if an extremal domain is homeomorphic to a ball, then it must be a ball; however Conjecture 6.2 is still open for domains whose boundary contains more than one component, or domains (such as a torus) that are not homeomorphic to a ball.

References

1. L. Ahlfors and A. Beurling, Conformal invariants and function theoretic null sets, Acta Math. **83** (1950), 101-129.
2. D. Aharonov and H.S. Shapiro, Domains in which analytic functions satisfy quadrature identities, J. Analyse Math. **30** (1976), 39-73.
3. H. Alexander, Projections of polynomial hulls, J. Funct. Anal. **3** (1973), 13-19.
4. C. Bandle, Isoperimetric Inequalities and Applications, Pitman, Boston-London-Melbourne, 1980.
5. B. Berge and J. Peseux, Variable focal lens controlled by an external voltage: An application of electrowetting, Eur. Phys. J. E **3** (2000), 159-163.
6. V. Blåsjö, The evolution of the isoperimetric problem, American Mathematical Monthly **112** (2005), no. 6, 526-566.

7.

33. On uniform approximation by harmonic functions, *Mich. Math. J.* **34**, 465-473, 1987.
34. D. Khavinson, Duality and uniform approximation by solutions of elliptic equations. *Contributions to operator theory and its applications (Mesa, AZ, 1987)*, 129-141, *Oper. Theory Adv. Appl.*, 35, Birkhuser, Basel, 1988.
35. D. Khavinson, An isoperimetric problem, *Linear and Complex Analysis, Problem Book 3, Part II*, edited by V. P. Khavin and N. K. Nikolski, *Lecture notes in Math.*, **1574** (1994), 133-135.
36. D. Khavinson, A. Solynin and D. Vassilev, Overdetermined boundary value problems, quadrature domains and applications, *Comput. Methods Funct. Theory* (5) 2005, No. 1, 19-48.
37. S. Ya. Khavinson, Two papers on extremal problems in complex analysis, *Amer. Math. Soc. Transl.* (2) **129** (1986).
38. S. Ya. Khavinson and G. Tumarkin, On the definition of analytic functions of class E^p in multiply-connected domains, *Uspehi Mat. Nauk* **13** (1958), no. 1 (79), 201-206 (in Russian).
- 39.

57. H. S. Shapiro, The Schwarz function and its generalizations to higher dimensions, John Wiley & Sons, 1992.
58. H. S. Shapiro, Remarks concerning domains of Smirnov type, Michigan Math. J. **13** (1966), 341-348.
59. J. Steiner, Einfache Beweise der isoperimetrischen Hauptsätze, J. reine ang. Math. **18** (1838), 281-296.
60. I. N. Vekua, Generalized Analytic Functions, Pergamon, London, 1962.
61. H. Weinberger, Remark on the preceding paper of Serrin, Arch. Rational Mech. Anal. **43** (1971), 319-320.

Department of Mathematics and Computer Science, Seton Hall University,
South Orange , New Jersey 07079

E-mail address: beneteca@shu.edu

Department of Mathematics, University of Arkansas, Fayetteville, Arkansas
72701

E-mail address: dmitry@uark.edu

Current address: 4201 Wilson Boulevard, NSF/MPS/DMS, Suite 1025, Arlington,
VA, 22230

E-mail address: dkhavins@nsf.gov