


An index to assess the health and benefits of the global ocean

Benjamin S. Halpern^{1,2}, Catherine Longo¹, Darren Hardy¹, Karen L. McLeod³, Jameal F. Samhouri⁴, Steven K. Katona⁵, Kristin Kleisner⁶, Sarah E. Lester^{7,8}, Jennifer O'Leary¹, Marla Ranelletti¹, Andrew A. Rosenberg⁵, Courtney Scarborough¹, Elizabeth R. Selig⁵, Benjamin D. Best⁹, Daniel R. Brumbaugh¹⁰, F. Stuart Chapin¹¹, Larry B. Crowder¹², Kendra L. Daly¹³, Scott C. Doney¹⁴, Cristiane Elfes^{15,16}, Michael J. Fogarty¹⁷, Steven D. Gaines⁸, Kelsey I. Jacobsen⁸, Leah Bunce Karrer⁵, Heather 9663255Leal7h259Bun18ce2t9b0344bHea1905.1.9768102.6708576.0566Tm14Tj0Hea988r866361.5999n18ce252r13umba17ugh



⁶Sea Around Us Project, Fisheries Centre, University of British Columbia, Vancouver, British Columbia V6T 1Z4, Canada.⁷Marine Science Institute, University of California, Santa Barbara, California 93106, USA. ⁸Bren School of Environmental Science and Management, University of California, Santa Barbara, California 93106, USA. ⁹Nicholas School of the Environment, Duke University, Durham, North Carolina 27708, USA.¹⁰Center for Biodiversity and Conservation, American Museum of Natural History, New York, New York 10024, USA. ¹¹Institute of Arctic Biology, University of Alaska, Fairbanks, Alaska 99775, USA. ¹²Center for Ocean Solutions and Hopkins Marine Station, Stanford University, Monterey, California 93940, USA.¹³College of Marine Science, University of South Florida, St Petersburg, Florida 33705, USA. ¹⁴Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA. ¹⁵Department of

framework to assess ocean health and motivate better data collection
to strengthen future iterations of the index.

Germany scored highly (index 73) because eight goals performed well (excepting food provision and tourism and recreation). Because of flatter distributions and greater range in values, whereas approach to scoring ocean health departs from a purely protectionist one that would aim to maintain natural systems with minimal human impact. The index credits sustainable non-extractive and extractive use (Table 31 and Supplementary Information). Tourism and recreation except in places where such uses are prohibited (for example, no-take reserves), as well as preservationist goals.

Third, the index allows transparent assessment of how societal biodiversity scores may seem surprisingly high, but this result values influence perspectives on ocean health. Although we weighted goals equally to avoid presuming societal values, we recognized that people value ocean benefits differently. To evaluate potential consequences of unequal weighting, we calculated index scores for 1000 simulations of unequal weighting, we calculated index scores for 1000 simulations of all species (see Supplementary Information). Diving deeper into the index, current status is the main driver of individual goal extractive use, extractive use, and strongly extractive use values (see Supplementary Information for further details). Resulting global biodiversity and carbon storage due to decreasing species status index scores ranged from 56 to 67 across value sets (Fig. 4 and Supplementary Table 30; country-level average maximum difference for tourism and recreation and coastal livelihoods and economies 7.16 0.2). For a few countries, most notably Romania, Russia, French Guiana, Micronesia and Denmark (Fig. 6 and Supplementary Information).

Countries with identical or similar scores provide examples of how changing weights created important differences, altering index scores by up to 27 (Supplementary Table 30). Monte Carlo simulations of thousands of possible value sets produced similar results (index 60.16 0.003 (s.e.); min 50; max 70; Supplementary Fig. 5 and Supplementary Table 30). The preservationist perspective produced the highest index score, primarily because extraction-based goals generally scored low whereas non-extractive goals scored higher. Because goal weights can influence index scores, it is critical to determine societal values (weights) before index calculation. Choosing a single unequal weighting scheme for this global analysis would not have been appropriate as these weights will vary by country, region and community.

Exploring the index

Variation among country-level index and individual goal scores offers novel insights into causes and consequences of different levels of ocean health (Fig. 5). Index scores had a largely unimodal distribution, which is expected in composite indices. No country scored above 86 and most scored below 70. Natural products, carbon

could significantly improve ocean health by addressing multiple goals. More effective and comprehensive protection of coastal areas and species, as is being pursued under the Convention on Biological Diversity Aichi Biodiversity Targets for 2020 (ref. 29), would directly benefit •sense of place• and •biodiversity• goals, and indirectly benefit most other goals by increasing ecological resilience and thus the likelihood of future goal delivery. Efforts to promote coastal livelihoods, environmentally sensitive urbanization of the coastal zone and improved sanitation infrastructure would improve •coastal livelihoods and economies•, •tourism and recreation•, and •clean water• goals. Simulating specific management scenarios could provide guidance on which actions would have the greatest impact.

Sustainability into the future

Sustainable delivery of each goal is foundational to our definition of a healthy ocean and approach to modelling the index. The status of many goals incorporates a penalty for pursuing a goal in a way that hampers its future delivery, whereas the •likely future state• augments scores for goals expected to improve in the near-term future (see Supplementary Information). About half of the goals are getting

Global-scale analyses are useful for global comparisons but tend to be locally imprecise because of inherent challenges in using available global data sets. Future finer-scale applications will allow full exploration of how to best use and refine the index. By calling attention to specific data layers (and gaps), the index can stimulate better measurements, more focused management and, hopefully, accelerate progress towards a healthier ocean.

Developing the index required many assumptions and compromises (see Supplementary Information); here we elaborate on three. First, we limited the index to ten constituent goals primarily for parsimony and ease of communication while maintaining a structure complementary to other ecosystem benefit typologies such as in the Millennium Ecosystem Assessment. We recognize that this structure significantly influences our results. Second, gaps existed in many data sets that we used, requiring proxies or models to fill those gaps (see Supplementary Information). For example, international arrivals data provide a modest proxy for coastal tourism (tourism and recreation goal) and undervalue the goal in nations with significant domestic tourism. Likewise, no global data exist for important stressors such as illegal fishing, habitat loss rates and point-source pollution. By identifying these data gaps, the index can help motivate future data collection. In other cases, we had to forgo better quality, region-specific data to maintain global consistency. Future iterations of the index, including those at finer geographic scales, can incorporate new data as available. Better data will in turn allow for construction of improved models that show greater fidelity to each goal's intent, but may also cause scores to chan

8eevcbu(eas)1213(n8e8iu(e-1.137tind3uis)-36k9hes)1is dats(se)-es,6(ter)-315(dats)-ecd irh-2-ec2(ocean.)JTJ 0.99

$$I \sim \frac{\sum_{i=1}^N a_i I_i}{P}$$

where a_i is the goal-specific weight ($a_i \in [0, 1]$; default is $a_i = 1/N$) (see Supplementary Information) and I_i is the average value of present and likely future status ($I_i = (x_i + x_{i,F})/2$, for each goal). The present status of goal i is its present status value x_i relative to a reference point $x_{i,r}$ uniquely chosen for each goal following guiding principles (see Supplementary Information and ref. 34), and rescaled 0...100. The likely future status is a function of present status x_i , recent (5 year) trend T_i , pressures p_i , and factors that promote resilience r_i , such that

$$x_{i,F} = \alpha I_i + \beta T_i + \gamma (p_i - r_i) x_i$$

where the discount rate $\alpha = 0$ and the weighting term $\beta = 0.67$, giving trend twice the importance of the difference between resilience and pressures in determining likely future state (see Supplementary Information). We tested the sensitivity of results to assumptions about α and β and found minimal differences for near-term timeframes (see Supplementary Information). Assessment of the likely future status captures whether the present status is likely to persist, improve or decline in the near-term future, based on current status and trends, and is therefore an indication rather than prediction of the near-term future. Ecological pressures fall into five broad categories, pollution, habitat destruction, species introductions, fishing and climate change, and are weighted equally to social pressures (such as poverty, political instability and corruption), with resilience measures such as international treaties and ecological resilience included when they address pressures relevant to a particular goal (see Supplementary Information). The inclusion of these factors ensures that the index is responsive to changes that are reflected more slowly in the current state.

and any associated references are available in the online version of the paper.

1 01 .
P 1 01 .

1. Halpern, B. S. et al. A global map of human impact on marine ecosystems. *Science* **31**, 948...952 (2008).
2. Jackson, J. B. C. et al. Historical overfishing and the recent collapse of coastal ecosystems. *Science* **293**, 629...637 (2001).
3. Pandolfi, J. M. et al. Global trajectories of the long-term decline of coral reef ecosystems. *Science* **301**, 955...958 (2003).
4. Center for International Earth Science Information Network. Socioeconomic Data and Applications Center (SEDAC) <http://sedac.ciesin.columbia.edu/> (2005).
5. Kareiva, P., Lalasz, R. & Marvier, M. Conservation in the Anthropocene: Beyond solitude and fragility. *Breakthrough J.* No. 2 29...37 (2011).
6. Kareiva, P., Watts, S., McDonald, R. & Boucher, T. Domesticated nature: shaping landscapes and ecosystems for human welfare. *Science* **31**, 1866...1869 (2007).
7. Rockström, J. et al. A safe operating space for humanity. *Nature* **461**, 472...475 (2009).
8. Rockström, J. et al. Planetary boundaries: exploring the safe operating space for humanity. *Ecol. Soc.* **14**, 32 (2009).
9. National Ocean Council. Draft National Ocean Policy Implementation Plan, http://www.whitehouse.gov/sites/default/files/microsites/ceq/national_ocean_policy_draft_implementation_plan_01-12-12.pdf (2012).
10. Administration of Barack H. Obama. Executive Order 13547: Stewardship of the ocean, our coasts, and the great lakes (2010).
11. McLeod, K. L. & Leslie, H. M. (eds). *Ecosystem-Based Management for the Oceans* (Island, 2009).
12. Munoz-Erickson, T. A., Aguilar-Gonzalez, B. & Sisk, T. D. Linking ecosystem health indicators and collaborative management: a systematic framework to evaluate ecological and social outcomes. *Ecol. Soc.* **12**, 6 (2007).
13. Rapport, D. J. et al. Ecosystem health: the concept, the ISEH, and the important tasks ahead. *Ecosyst. Health* **2**, 82...90 (1999).
14. Cairns, J., McCormick, P. V. & Niederlehner, B. R. A proposed framework for developing indicators of ecosystem health. *Hydrobiologia* **261**, 1...44 (1993).
15. Kelly, J. R. & Harwell, M. A. Indicators of ecosystem recovery. *Environ. Manage.* **14**, 527...545 (1990).
16. Rapport, D. J., Costs.

▲ We measured ocean health as a function of ten widely held public goals (see Supplementary Information for further details) for what the ocean can provide

have specific consequences for goals, we assessed and ranked separately each ecological stressor within these categories.

To account for the cumulative effect of stressors, we summed the weighted intensities of each stressor within a pressure category and divided by the maximum weighted intensity that could be achieved by the worst stressor (max 3.0) such that:

$$p_k = \frac{\sum_i w_i s_i}{3}$$

where w_i is the stressor-specific sensitivity weights (from Supplementary Table 25) and s_i is the data-derived intensity of the associated stressor (which is scaled 0...1). If $s_i > 100$, we set the value equal to 100. This formulation assumes