GUIDELINES ON ANESTHESIA AND ANALGESIA IN LABORATORY ANIMALS

University of South Florida provides the following guidelines for use by IACUC-certified faculty and staff.

CONTENTS **PAGE**

A. Background…… 1
B. Definitions..…….. 2
C. General Considerations…….,…….. 3
D. Controlled Substances…… 3
E. Pre-Anesthetic Treatments…… 4
F. General Anesthetics…… 4
G. Neuromuscular Blocking Agents…….. 5
H. Monitoring Anesthesia…… 6
I. Analgesics…… 7
J. Comments regarding Anesthetics and Analgesics……... 7

REFERENCE TABLES **PAGE**

I. Signs of Pain and Distress in Laboratory Animals…… 10
II. Commonly Used Anesthetics and Analgesics for Mice…… 11
III. Commonly Used Anesthetics and Analgesics for Rats…… 12
IV. Commonly Used Anesthetics and Analgesics for Gerbils…….. 13
V. Commonly Used Anesthetics and Analgesics for Hamsters…… 14
VI. Commonly Used Anesthetics and Analgesics for Guinea Pigs…… 15
VII. Commonly Used Anesthetics and Analgesics for Rabbits…… 16
VIII. Commonly Used Anesthetics and Analgesics for Dogs…… 17
IX. Commonly Used Anesthetics and Analgesics for Cats…… 18
X. Commonly Used Anesthetics and Analgesics for Pigs ……..19
XI. Commonly Used Anesthetics and Analgesics for Sheep/Goats..……..20
XII. Commonly Used Anesthetics and Analgesics for Macaques ……...21
XIII. Commonly Used Anesthetics and Analgesics for Birds……22

A. Background

1. A fundamental responsibility of individuals that use animals in research, teaching or testing is to anticipate and eliminate or minimize any potential that procedures may cause animal pain, distress, or discomfort.

2. Although animals that are in pain may not behave like humans, (e.g., pain in animals may be accompanied by immobility and silence, in contrast to the groans and cries of human patients), it is assumed that **procedures that cause pain in humans cause pain in animals**.

3. The presence of pain in animals can be recognized by alterations in animal behavior (e.g., reduced activity, reduced grooming, hunched-up posture, altered gait, changes in temperament, vocalizations, reduced food and water intake, reduced urinary and fecal output), and in physiological variables, (e.g., reduced depth of respiration, increased heart rate, and reduced hydration status) (refer to Table I).

4. Animal pain, distress, and discomfort can **produce a range of undesirable h/ledT1(h/ḃ)123ḃ8 ḃḃ.6c}8 (T/ledT1ḃoo8 (T**

- **Schedule V** - A drug or other substance that has a low potential for abuse relative to the drugs or substances in schedule IV, and has a currently accepted medical use in treatment in the United States. Abuse of the drug or other substance may lead to limited physical dependence or psychological dependence relative to the drug or other substance in schedule IV.

E. Pre-Anesthetic Treatments

1. Pigs, cats, dogs, and primates should be fasted for 8-12 hours to minimize the risk of vomiting and

their staff are trained in the proper use of tranquilizers, anesthetics, and analgesics appropriate for the species and planned procedures.

2. Many factors can affect the activity of anesthetics. The species, strain, sex, age, nutritional and disease status, relative body size, disposition/demeanor, presence of concurrent pain or distress, or medication are known to cause a variation in the amount of drug needed to produce a desired effect in an individual animal.

3. Although the mechanisms of action vary, anesthetics produce, in a controllable manner, both loss of consciousness and an absence of motor response to noxious stimuli. This unconsciousness, analgesia, and muscle relaxation should be sufficient to allow the performance of procedures without the subject experiencing pain. In addition to these effects, anesthetics also produce a depressant effect on the cardiovascular, respiratory, and thermo-regulatory systems. Their use must be monitored closely.

4. The level of anesthesia should be limited to the induction of the minimal degree of CNS depression necessary for performing the procedure. When an injected anesthetic agent is used, drug dose calculation should be based on body weight and age. General anesthesia must be given "to effect," as noted in physiologic responses and in response to noxious stimuli. It is important to realize that some drugs take time to take effect. Anesthetic death can be attributed to giving the anesthetic insufficient time to work. This is especially true of parentally administered drugs (e.g., barbiturates). Once they are injected, there is little the anesthetist can do to control the outcome.

5. **Inhaled Anesthetics** (e.g., halothane, enflurane, isoflurane, sevoflurane, desflurane) have a greater margin of safety and produce a more stable plane of surgical anesthesia, when used with a calibrated vaporizer, than injectable anesthetics. Since these anesthetics ente5.and leav 5bo (c)-8.1.1 (h)-12.2 (e)-12.3 (y)1v(an)-

for days after administration. Therefore, it is important to check animals for signs of anorexia, fever, vomiting, or abnormal respiration or heart rate.

7. Indications of Anesthetic Overdose – Monitoring vital signs continuously during anesthesia will provide early warning of potential problems and emergencies that may be averted by appropriate and quick corrective actions. Do not rely on a single parameter to assess the animal's condition. All parameters should be evaluated prior to initiating any corrective actions. The following indicators of anesthetic overdose, which may lead to cardiac or respiratory failure, are helpful in assessing the animal's status during anesthesia. Heart rate may be rapid or slow, depending on the animal's state of physiological decompensation. Remember that anticholinergics cause the heart rate to increase. Pulses may be

J. Comments Regarding Anesthetics and Analgesics

1. Several commonly used or historically used anesthetics and analgesic medications are described briefly below. However, numerous additional agents are available for use in a variety of species. Contact a University of South Florida laboratory animal veterinarian for additional information on drugs not listed here. A veterinary drug formulary and a number of veterinary anesthesia textbooks are available in the Comparative Medicine library.

2. **Acepromazine Maleate** (formerly acetylpromazine), a phenothiazine derivative, is a potent neuroleptic agent with relatively low toxicity. Acepromazine induces tranquilization, muscle relaxation, and a decrease in spontaneous activity. At high doses, sedation occurs. Preanesthetic administration decreases the amount of general anesthetic required. Acepromazine possesses antiemetic, anticonvulsant, antispasmodic, hypotensive, and hypothermic properties. Acepromazine will prevent or decrease severity of the malignant hyperthermia syndrome in susceptible swine exposed to halothane. Acepromazine potentiates opiates such as butorphanol and buprenorphine, which if used in combination as a pre-anesthetic, will provide sedation as well as preemptive analgesia.

3. **Fentanyl citrate patch** is a very potent opiate agonist. The patch is a transdermal delivery system for the fentanyl, and is used primarily in pigs, dogs, and cats to alleviate postoperative pain, and to control chronic pain (e.g. associated with cancer). Therapeutic levels are achieved within 6-8 hours of application in the cat, while it takes at least 12 hours to reach therapeutic levels in the dog, so patch application should be performed prior to the procedure keeping these times in mind. Small dogs and

should be used for these anesthetic agents because lethal concentrations can easily be reached using the open drop method, or using a bell jar as an anesthetic chamber.

10. **Xylazine** (i.e., Rompun®) is a centrally acting alpha-

Table I. Signs of Pain and Distress in Laboratory Animals*

Species	Signs of mild to moderate pain of distress	Signs of severe or chronic pain or distress
Mouse	Eyelids partially closed; changes in respiration; rough hair coat; increased vibrissae movement; unusually apprehensive or aggressive; possible writhing, scratching, biting, self-mutilation; hunched posture; sudden running; aggressive vocalization; guarding.	Weight loss; dehydration; incontinence; soiled hair coat; eyes sunken, lids closed; wasting of muscles on back; sunken or distended abdomen; decreased vibrissae movement; unresponsive; separates from group; hunched posture; ataxia; circling; hypothermia; decreased vocalization.

duplicate/grd/epost/1 pp 6 17 (1)10 7 (1)8 (1)8 (1)8 (s)-1 (e)3 (0d ()2 .1 (ur)3.6 (e0.

Table II. Commonly Used Anesthetics and Analgesics for Mice

University of South Florida provides the following table as a reference only, for use by IACUC-certified faculty and staff only.

C06.44 5f5076 7252.48 20 Td 2.88r 3n5BTf*53.2 r T9.[C17.64 5f507f*506.44 20 Td 13.[C 3n5BTTf506.44 T9.[C113.[C . 3 nBTf506]Do4

Anesthesia in Mice	Dose & Route

Table III. Commonly Used Anesthetics and Analgesics for Rats

University of South Florida provides the following table as a reference only, for use by IACUC

Table IV. Commonly Used Anesthetics and Analgesics for Gerbils

University of South Florida provides the following table as a reference only, for use by IACUC-certified faculty and staff only.

Anesthesia in Gerbils	Dose & Route	Comments
Isoflurane (Forane®)	To effect. In general, 1-4%	Precision vaporizer, adequate ventilation or scavenging essential K9 044 2396 13.3
Ketamine + Xylazine	50 mg/kg (K) + 2 mg/kg (X) IP	Duration 20-50min

Table IX. Commonly Used Anesthetics and Analgesics for Cats

University of South Florida provides the following table as a reference only, for use by IACUC-certified faculty and staff only.

Anesthesia in Cats	Dose & Route	Comments
Isoflurane (Forane®	To effect. In general, 3-4% induction, 1-2% maintenance; inhalation	Precision vaporizer, adequate ventilation or scavenging essential

Table XI. Commonly Used Anesthetics and Analgesics for Sheep and Goats

University of South Florida provides the following table as a reference only, for use by IACUC-certified faculty and staff only.

Anesthesia in Sheep & Goats	Dose & Route	Comments
[redacted]	To effect. In general, 3-4% induction, 1-	Precision vaporizer, adequate ventilation or scavenging essential
Diazepam + Ketamine	0.3 – 0.5 mg/kg (D) + 5 mg/kg (K) IV, IM	Up to 20 minutes of anesthesia; for minor procedures
Xylazine + Ketamine	0.05 - 1	

Table XIII. Commonly Used Anesthetics and Analgesics for Birds

jEMC